



Anaphylaxis Policy

Purpose:

To explain to Whittlesea Secondary College parents, carers, staff and students the processes and procedures in place to support students diagnosed as being at risk of suffering from anaphylaxis. This policy also ensures that Whittlesea Secondary College is compliant with Ministerial Order 706 and the Department's guidelines for anaphylaxis management.

Scope:

This Policy applies to:

- all Whittlesea Secondary College staff, including casual relief staff and volunteers
- all Whittlesea Secondary College students who have been diagnosed with anaphylaxis, or who may require emergency treatment for an anaphylactic reaction, and their parents and carers.

Policy:

School Statement

Whittlesea Secondary College will fully comply with Ministerial Order 706 and the associated guidelines published by the Department of Education.

Anaphylaxis

Anaphylaxis is a severe allergic reaction that occurs after exposure to an allergen. The most common allergens for school-aged children are nuts, eggs, cow's milk, fish, shellfish, wheat, soy, sesame, latex, certain insect stings and medication.

Symptoms

Signs and symptoms of a mild to moderate allergic reaction can include:

- swelling of the lips, face and eyes
- hives or welts
- tingling in the mouth.

Signs and symptoms of anaphylaxis, a severe allergic reaction, can include:

- difficult/noisy breathing
- swelling of tongue
- difficulty talking and/or hoarse voice
- wheeze or persistent cough
- persistent dizziness or collapse
- student appears pale or floppy



- abdominal pain and/or vomiting.

Symptoms usually develop within ten minutes and up to two hours after exposure to an allergen but can appear within a few minutes.

Treatment

Adrenaline given as an injection into the muscle of the outer mid-thigh is the first aid treatment for anaphylaxis.

Individuals diagnosed as being at risk of anaphylaxis are prescribed an adrenaline autoinjector for use in an emergency. These adrenaline autoinjectors are designed so that anyone can use them in an emergency.

Individual Anaphylaxis Management Plans

All students at Whittlesea Secondary College who are diagnosed by a medical practitioner as being at risk of suffering from an anaphylactic reaction **MUST** have an **Individual Anaphylaxis Management Plan**.

When notified of an anaphylaxis diagnosis, the Principal of Whittlesea Secondary College is responsible for developing a plan in consultation with the student's parents/carers.

Where necessary, an Individual Anaphylaxis Management Plan will be in place as soon as practicable after a student enrolls at Whittlesea Secondary College and where possible, before the student's first day.

Parents and carers must:

- obtain an ASCIA Action Plan for Anaphylaxis from the student's medical practitioner and provide a copy to the College as soon as practicable.
- immediately inform the College in writing if there is a relevant change in the student's medical condition and obtain an updated ASCIA Action Plan for Anaphylaxis.(RED)
- provide an up-to-date photo of the student for the ASCIA Action Plan for Anaphylaxis (RED) when that Plan is provided to the College and each time it is reviewed.
- provide the College with a current adrenaline autoinjector for the student that has not expired.
- participate in annual reviews of the student's Plan.

Each student's Individual Anaphylaxis Management Plan must include:

- information about the student's medical condition that relates to allergies and the potential for anaphylactic reaction, including the type of allergies the student has.
- information about the signs or symptoms the student might exhibit in the event of an allergic reaction based on a written diagnosis from a medical practitioner.



- strategies to minimise the risk of exposure to known allergens while the student is under the care or supervision of College staff, including in the school yard, at camps and excursions, or at special events conducted, organised or attended by the College.
- the name of the person(s) responsible for implementing the risk minimisation strategies, which have been identified in the Plan.
- information about where the student's medication will be stored.
- the student's emergency contact details.
- an up-to-date ASCIA Action Plan for Anaphylaxis (RED) completed by the student's medical practitioner.

Review and updates to Individual Anaphylaxis Management Plans

A student's Individual Anaphylaxis Management Plan will be reviewed and updated on an annual basis in consultation with the student's parents/carers. The plan will also be reviewed and, where necessary, updated in the following circumstances:

- as soon as practicable after the student has an anaphylactic reaction at the College
- if the student's medical condition, insofar as it relates to allergy and the potential for anaphylactic reaction, changes
- when the student is participating in an off-site activity, including camps and excursions, or at special events including fetes and concerts.

Whittlesea Secondary College may also consider updating a student's Individual Anaphylaxis Management Plan if there is an identified and significant increase in the student's potential risk of exposure to allergens at school.

Location of plans and adrenaline autoinjectors

- A copy of each student's Individual Anaphylaxis Management Plan will be stored with their ASCIA Action Plan for Anaphylaxis at the School Office and First Aid Room. These medications will be labelled with the student's name and photo and stored in a College-supplied RED Medication pouch.
- Some Students may also choose to keep an additional adrenaline autoinjector on their person. They should ensure they keep a copy of their most recent ASCIA Anaphylaxis Action Plan (RED) together with their medication.



Risk Minimisation Strategies

To reduce the risk of a student suffering from an anaphylactic reaction at Whittlesea Secondary College, we have put in place the following strategies:

- College staff and students are regularly reminded to wash their hands after eating;
- College students are discouraged from sharing food
- Garbage bins at the College are to remain covered with lids to reduce the risk of attracting insects
- Gloves must be worn when picking up papers or rubbish in the College grounds
- College canteen staff are trained in appropriate food handling to reduce the risk of cross-contamination
- A general use adrenaline autoinjector will be stored at the First Aid Room
- Planning for off-site activities will include risk minimisation strategies for students at risk of anaphylaxis including supervision requirements, appropriate number of trained staff, emergency response procedures and other risk controls appropriate to the activity and students attending.

Adrenaline autoinjectors for general use

Whittlesea Secondary College will maintain a supply of adrenaline autoinjectors for general use, as a back-up to those provided by parents and carers for specific students, and also for students who may suffer from a first-time reaction at the College.

There are currently **4 adrenaline devices** approved by the Therapeutic Goods Administration for use in Australia:

1. **the EpiPen®**
2. **the Anapen®**
3. **Jext®**
4. **Neffy®.**

All devices can be used when provided by families for students, however, the principal or allocated staff member can only use EpiPen®, Anapen® or Jext® adrenaline autoinjector for general use. For more information about which autoinjector to purchase for general use, refer to [Adrenaline autoinjectors for general use](#).

Adrenaline autoinjectors for general use will be stored at:

- The Main Office
- Science rooms
- PITT House (Senior School)
- Cooking Rooms



- Trade Training Centre
- Gym
- Wellbeing Centre
- Trauma Bag in Main Office

These Adrenaline autoinjectors will be labelled “general use”.

The Principal is responsible for arranging the purchase of adrenaline autoinjectors for general use, and will consider:

- the number of students enrolled at Whittlesea Secondary College at risk of anaphylaxis.
- the accessibility of adrenaline autoinjectors supplied by parents.
- the availability of a sufficient supply of autoinjectors for general use in different locations at the College, as well as at camps, excursions and events.
- the limited life span of adrenaline autoinjectors, and the need for general use adrenaline autoinjectors to be replaced when used or prior to expiry.
- the weight of the students at risk of anaphylaxis to determine the correct dosage of adrenaline autoinjector/s to purchase.

Emergency Response

In the event of an anaphylactic reaction, the emergency response procedures in this policy must be followed, together with the Colleges general first aid procedures, emergency response procedures and the student’s Individual Anaphylaxis Management Plan.

A complete and up-to-date list of students identified as being at risk of anaphylaxis is maintained by the First Aid Officer and stored at the First Aid Office.

For camps, excursions and special events, a designated staff member will be responsible for maintaining a list of students at risk of anaphylaxis attending the special event, together with their Individual Anaphylaxis Management Plans and adrenaline autoinjectors, where appropriate.

If a student experiences an anaphylactic reaction at school or during a school activity, school staff must:

STEP	ACTION
1.	<ul style="list-style-type: none"> • Lay the person flat • Do not allow them to stand or walk • If breathing is difficult, allow them to sit



	<ul style="list-style-type: none">• Be calm and reassuring• Do not leave them alone• Seek assistance from another staff member or reliable student to locate the student's adrenaline autoinjector or the school's general use autoinjector, and the student's Individual Anaphylaxis Management Plan, stored at the First Aid Office.• If the student's plan is not immediately available, or they appear to be experiencing a first-time reaction, follow steps 2 to 5
2.	<p><u>Administer an EpiPen or EpiPen Jr</u></p> <ul style="list-style-type: none">• Remove from plastic container• Form a fist around the EpiPen and pull off the blue safety release (cap)• Hold leg still and place orange end against the student's outer mid-thigh (with or without clothing)• Push down hard until a click is heard or felt and hold in place for 3 seconds• Remove EpiPen• Note the time the EpiPen is administered• Retain the used EpiPen to be handed to ambulance paramedics along with the time of administration. <p>OR</p> <p><u>Administer an Anapen® 500</u></p> <ul style="list-style-type: none">• Pull off the black needle shield• Pull off grey safety cap (from the red button)• Place needle end firmly against the student's outer mid-thigh at 90 degrees (with or without clothing)• Press red button so it clicks and hold for 3 seconds• Remove Anapen®• Note the time the Anapen is administered• Retain the used Anapen to be handed to ambulance paramedics along with the time of administration. <p>OR</p> <p><u>Administer Jext 150 or 300</u></p> <ul style="list-style-type: none">• Form fist around Jext and pull off yellow cap• Place black injector tip against outer-mid thigh (with or without clothing)• Push black tip firmly until a click is heard and hold in place for 3 seconds.• Remove Jext• Note the time the Jext device is administered.• The used adrenaline device must be handed to the ambulance paramedics along with the time of administration



	<p>OR</p> <p><u>Administer Neffy® 1mg or 2mg</u></p> <ul style="list-style-type: none">• Hold the nasal spray with your thumb on the bottom of the plunger and a finger on either side of the nozzle.• Do not pull or push on the plunger.• Do not test or prime (pre-spray).• Each Neffy nasal spray contains only one spray.• Place the nozzle of the nasal spray into a nostril until fingers touch the nose.• For smaller nostrils, aim for the fingers to touch the nose.• Keep the nozzle pointed towards the forehead. Do not angle the nozzle of the nasal spray to the inner or outer walls of the nose.• Press the plunger up firmly until the dose is administered and it sprays into the nostril. Note the time the Neffy device is administered.• The used adrenaline device must be handed to the ambulance paramedics along with the time of administration
3.	Call an ambulance (000)
4.	If there is no improvement or severe symptoms progress (as described in the ASCIA Action Plan for Anaphylaxis(RED)), further adrenaline doses may be administered every five minutes, if other adrenaline autoinjectors are available.
5.	Contact the student's emergency contacts.
6.	The principal or a staff member allocated to do so must contact the Incident Support and Operations Centre (ISOC) on 1800 126 126 to report 'High' or 'Extreme' severity incidents to report the incident. Incidents assessed as 'Low' or 'Medium' can be reported directly into EduSafe Plus by the principal or their allocated staff member.



If a student appears to be having a severe allergic reaction but has not been previously diagnosed with an allergy or being at risk of anaphylaxis, school staff should follow steps 2 – 5 as above.

For first time anaphylactic reactions, the school's general use adrenaline autoinjector device must be used. If the general use device is not immediately available in an anaphylaxis emergency, College staff may use another student's adrenaline device, including the Epipen®, Anapen®, Jext® or Neffy® device. This may save a life. If another student's adrenaline device is used in an anaphylaxis emergency, the College must notify the parents of the student whose device was used and immediately replace the device.

Where possible, schools should consider using the correct dose of adrenaline autoinjector depending on the weight of the student. However, in an emergency if there is no other option available, any device should be administered to the student.

Communication Plan:

This policy will be available on Whittlesea Secondary College's Website so that parents and other members of the College community can easily access information about Whittlesea Secondary College's anaphylaxis management procedures. The parents and carers of students who are enrolled at Whittlesea Secondary College and are identified as being at risk of anaphylaxis will also be provided with a copy of this policy.

The Principal is responsible for ensuring that all relevant College staff, including casual relief staff, canteen staff and volunteers are aware of this policy and Whittlesea Secondary College's procedures for anaphylaxis management.

Casual relief staff and volunteers who are responsible for the care and/or supervision of students who are identified as being at risk of anaphylaxis will also receive a verbal briefing on this policy, their role in responding to an anaphylactic reaction and where required, the identity of students at risk.

The Principal is also responsible for ensuring relevant College staff are trained and briefed in anaphylaxis management, consistent with the Department's [Anaphylaxis Guidelines](#).

Staff training

The Principal will ensure that the following College staff are appropriately trained in anaphylaxis management:

- College staff who conduct classes attended by students who are at risk of anaphylaxis
- College staff who conduct specialist classes, canteen staff, administration staff, first aiders and any other member of College staff as required by the Principal based on a risk assessment.

Staff who are required to undertake training must have completed:

- an approved face-to-face anaphylaxis management training course in the last three years, or
- an approved online anaphylaxis management training course in the last year.



Whittlesea Secondary College uses the following training course:

[ASCIA eTraining Course](#) with 22579VIC and 22578VIC.

Staff are also required to attend a briefing on anaphylaxis management and this policy at least twice per year (with the first briefing to be held at the beginning of the school year), facilitated by the nominated School anaphylaxis Supervisor/s who has successfully completed an anaphylaxis management course within the last 2 years, and familiarise themselves with locations of items. (See Appendix 1 Map)

Each Briefing will address:

- The College's anaphylaxis management policy
- The causes, symptoms and treatment of anaphylaxis
- The identities of students with a medical condition that relates to allergies and the potential for anaphylactic reaction, and where their medication is located
- How to use an adrenaline autoinjector, including hands on practice with a trainer adrenaline autoinjector
- The College's general first aid and emergency response procedures
- The location of, and access to, adrenaline autoinjectors that have been provided by parents or purchased by the College for general use.
- How to access on-going support and training.

When a new student enrolls at Whittlesea Secondary College who is at risk of anaphylaxis, the College will develop an interim plan in consultation with the student's parents and ensure that appropriate College staff are trained and briefed as soon as possible.

A record of College staff training courses and briefings will be maintained through the College's online Emergency Management Plan.

T

he Principal will ensure that while students at risk of anaphylaxis are under the care or supervision of the College outside of normal class activities, including in the College yard, at camps and excursions, or at special event days, there is enough College staff present who have been trained in anaphylaxis management.



Further Information and Resources:

This policy should be read in conjunction with the following College and Department policies and guidelines:

- [Anaphylaxis](#)
- [Allergies](#)
- [First Aid for Students and Staff](#)
- [Health Care Needs](#)
- [Managing Reporting Incidents \(including Emergencies\)](#)
- [Medication](#)

Other related Resources:

- [Allergy & Anaphylaxis Australia](#)
- ASCIA Guidelines: [Schooling and childcare](#)
- Royal Children’s Hospital: [Allergy and immunology](#)
- [Hero HQ Training](#)
- [Whittlesea Secondary College Medications Policy](#)
- [Whittlesea Secondary College Duty of Care Policy](#)
- [Whittlesea Secondary College First Aid Policy](#)

Policy Review and Approval:

The Principal will complete the Departments Annual Risk Management Checklist for anaphylaxis Management to assist with the evaluation and review of this policy and the support provided to students at risk of Anaphylaxis.

Policy last reviewed	February 2026
Consultation	First Aid Officers, Anaphylaxis Supervisor, Principal Executive
Approval by	Principal
Next scheduled review date	February 2028



Appendix 1

Whittlesea Secondary College

